



Phycoremediation potential of brown macroalgae species *Saccharina latissimi* and *Laminaria digitata* towards inorganic arsenic in a multitrophic pilot-scale experiment

Rasmussen, Rie Romme; Cunha, Sara; Fernandes, José; Oliveira, Cátia; Anacleto, Patrícia; Barbosa, Vera; Alves, Ricardo N.; Marques, António T. ; H.M. van den Heuvel, Freek; Sloth, Jens Jørgen

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Rasmussen, R. R., Cunha, S., Fernandes, J., Oliveira, C., Anacleto, P., Barbosa, V., Alves, R. N., Marques, A. T., H.M. van den Heuvel, F., & Sloth, J. J. (2017). *Phycoremediation potential of brown macroalgae species *Saccharina latissimi* and *Laminaria digitata* towards inorganic arsenic in a multitrophic pilot-scale experiment*. Poster session presented at Seafood Safety, Brussels, Belgium.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Phycoremediation potential of brown macroalgae species *Saccharina latissima* and *Laminaria digitata* towards inorganic arsenic in a multitrophic pilot-scale experiment

Rasmussen RR¹; Cunha S²; Fernandes J²; Oliveira C²; Anacleto P³;
Barbosa V³; Alves R³; Marques A³; van den Heuvel F⁴; Sloth JJ¹

1) Technical University of Denmark (DTU), National Food Institute, Søborg, Denmark
2) LAQV- REQUIMTE, Laboratory of Bromatology and Hydrology, Faculty of Pharmacy, University of Porto, Portugal 3) Portuguese Institute for the Sea and Atmosphere (IPMA), Lisboa, Portugal
4) Hortimare Projects & Consultancy BV, The Netherlands



Intro. Phycoremediation has emerged as one of the most promising and cost-effective technologies, which uses the macroalgae capacity to remove, degrade or render harmful contaminants in aquatic systems. Arsenic (As) is a ubiquitous metalloid which is bioaccumulated in the marine food chain and inorganic arsenic (iAs) is considered to be the most toxic form of arsenic.

Aim. To evaluate the phycoremediation capacity of the two brown seaweed species Sugar kelp (*Saccharina latissima*, SL) and Oarweed (*Laminaria digitata*, LD) in a controlled multitrophic (water, algae, mussels) pilot experiment with 0, 6, 12 h, ... up to 15 days (360 h) exposure to iAs.

Results.




Figure 1. As conc. (mean±SD) in A) seawater sampled twice and B) seaweed from tanks added iAs with dead or alive LD and SL.

- Water (As_{total}) LD < SD. Dead algae releases As. * (SL) biomass released As.
- Seaweed (iAs) LD > SD in absolute levels. Low iAs in dead biomass.

Conclusion. *Laminaria digitata* was most efficient for removal of arsenic from seawater and hence a better choice for phycoremediation practices towards this parameter.




Figure 3. As speciation for LD, 12 h exposure, 20 µg iAs/l. • iAs << As_{total}